

## University of Groningen

### Size matters

Eklöf, Johan; Austin, Åsa; Bergström, Ulf; Donadi, Serena; Eriksson, Britas D H K; Hansen, Joakim; Sundblad, Göran

*Published in:*  
PeerJ

*DOI:*  
[10.7717/peerj.2906](https://doi.org/10.7717/peerj.2906)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2017

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Eklöf, J., Austin, Å., Bergström, U., Donadi, S., Eriksson, B. D. H. K., Hansen, J., & Sundblad, G. (2017). Size matters: Relationships between body size and body mass of common coastal, aquatic invertebrates in the Baltic Sea. *PeerJ*, 5, [e2906]. <https://doi.org/10.7717/peerj.2906>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Size matters: relationships between body size and body mass of common coastal, aquatic invertebrates in the Baltic Sea

Johan Eklöf<sup>1</sup>, Åsa Austin<sup>1</sup>, Ulf Bergström<sup>2</sup>, Serena Donadi<sup>1,3</sup>, Britas D.H.K. Eriksson<sup>4</sup>, Joakim Hansen<sup>3</sup> and Göran Sundblad<sup>5</sup>

<sup>1</sup> Department of Ecology, Environment and Plant Sciences (DEEP), Stockholm University, Stockholm, Sweden

<sup>2</sup> Department of Aquatic Resources, Swedish University of Agricultural Sciences, Öregrund, Sweden

<sup>3</sup> Baltic Sea Centre, Stockholm University, Stockholm, Sweden

<sup>4</sup> Groningen Institute for Evolutionary Life-Sciences GELIFES, University of Groningen, Groningen, Netherlands

<sup>5</sup> Aquabiota Water Research, Stockholm, Sweden

## ABSTRACT

**Background.** Organism biomass is one of the most important variables in ecological studies, making biomass estimations one of the most common laboratory tasks. Biomass of small macroinvertebrates is usually estimated as dry mass or ash-free dry mass (hereafter ‘DM’ vs. ‘AFDM’) per sample; a laborious and time consuming process, that often can be speeded up using easily measured and reliable proxy variables like body size or wet (fresh) mass. Another common way of estimating AFDM (one of the most accurate but also time-consuming estimates of biologically active tissue mass) is the use of AFDM/DM ratios as conversion factors. So far, however, these ratios typically ignore the possibility that the relative mass of biologically active vs. non-active support tissue (e.g., protective exoskeleton or shell)—and therefore, also AFDM/DM ratios—may change with body size, as previously shown for taxa like spiders, vertebrates and trees.

**Methods.** We collected aquatic, epibenthic macroinvertebrates (>1 mm) in 32 shallow bays along a 360 km stretch of the Swedish coast along the Baltic Sea; one of the largest brackish water bodies on Earth. We then estimated statistical relationships between the body size (length or height in mm), body dry mass and ash-free dry mass for 14 of the most common taxa; five gastropods, three bivalves, three crustaceans and three insect larvae. Finally, we statistically estimated the potential influence of body size on the AFDM/DM ratio per taxon.

**Results.** For most taxa, non-linear regression models describing the power relationship between body size and (i) DM and (ii) AFDM fit the data well (as indicated by low SE and high  $R^2$ ). Moreover, for more than half of the taxa studied (including the vast majority of the shelled molluscs), body size had a negative influence on organism AFDM/DM ratios.

**Discussion.** The good fit of the modelled power relationships suggests that the constants reported here can be used to quickly estimate organism dry- and ash-free dry mass based on body size, thereby freeing up considerable work resources. However, the considerable differences in constants between taxa emphasize the need for taxon-specific relationships, and the potential dangers associated with ignoring body size. The negative influence of body size on the AFDM/DM ratio found in a majority of the molluscs could be caused by increasingly thicker shells with organism age, and/or

Submitted 25 August 2016  
Accepted 12 December 2016  
Published 25 January 2017

Corresponding author  
Johan Eklöf, johan.eklof@su.se

Academic editor  
Richard Taylor

Additional Information and  
Declarations can be found on  
page 11

DOI 10.7717/peerj.2906

© Copyright  
2017 Eklöf et al.

Distributed under  
Creative Commons CC-BY 4.0

## OPEN ACCESS

spawning-induced loss of biologically active tissue in adults. Consequently, future studies utilizing AFDM/DM (and presumably also AFDM/wet mass) ratios should carefully assess the potential influence of body size to ensure more reliable estimates of organism body mass.

**Subjects** Aquaculture, Fisheries and Fish Science, Biodiversity, Ecology, Marine Biology, Zoology  
**Keywords** Estuary, Biometry, Infauna, Submerged aquatic vegetation, Isometric scaling, Length:weight relationship, Epifauna, Allometry, Seagrass, Weight

## INTRODUCTION

Organism biomass is inarguably one of the more important variables in ecology, playing a central role in studies ranging from ecophysiology and community and food web regulation, to whole-ecosystem metabolism (e.g., [Enquist & Niklas, 2001](#); [Gruner et al., 2008](#); [Perez-Harguindeguy et al., 2013](#)). As a consequence, to accurately estimate organism biomass constitutes one of the most common and important tasks in ecological studies.

Small invertebrates retained on 0.5–1 mm sieves (hereafter ‘macrofauna’) make up a major part of animal density, diversity and biomass in many ecosystems; e.g., insects and arachnids in terrestrial ecosystems; epibenthic, aquatic crustaceans, echinoderms and molluscs in stands of aquatic vegetation; and infaunal (sediment-dwelling) worms, crustaceans and molluscs in marine sediments. Macrofauna biomass is typically reported as dry- or ash-free dry mass per unit area (e.g., g per m<sup>2</sup>), which requires observers to repeatedly identify, sort, dry and weigh individual or pooled organisms; a time-consuming, expensive and tedious process. Many studies have shown that more easily measured proxy variables scale predictably with dry mass and therefore can be used to speed up biomass estimations; e.g., wet (fresh) mass ([Brey, Rumohr & Ankar, 1988](#); [Ricciardi & Bourget, 1998](#); [Brey et al., 2010](#)) and body size, based on either exact length measurement ([Smock, 1980](#); [Frithsen, Rudnick & Doering, 1986](#); [Sabo, Bastow & Power, 2002](#)) or retention on sieves of certain mesh sizes ([Widbom, 1984](#); [Edgar, 1990](#); [Casagrande & Boudouresque, 2002](#)). While wet mass can be a very good proxy (see e.g., [Ricciardi & Bourget, 1998](#)), we—as others before—argue that body size (e.g., length) holds several advantages, and lacks several disadvantages associated with wet mass estimations. First, ecological theory supported by empirical data suggest body mass scales predictably with length in the form of power relations ([Smock, 1980](#); [Sabo, Bastow & Power, 2002](#)). Second, while freezing/thawing and fixation in conservation liquids (e.g., EtOH or formalin) can affect both organism wet mass ([Howmiller, 1972](#); [Mason, Lewis & Weber, 1983](#); [Leuven, Brock & van Druten, 1985](#)) and length ([Hjörleifsson & Klein-MacPhee, 1992](#); [Kapiris, Miliou & Moraitou-Apostolopoulou, 1997](#)), wet mass estimations are also very sensitive to exactly how specimens are blotted, centrifuged (to remove excess water), and exposed to air and light before and during weighing ([Howmiller, 1972](#); [Mason, Lewis & Weber, 1983](#); [Leuven, Brock & van Druten, 1985](#)). As size estimations do not require blotting, they are less sensitive to observer error, and also faster to perform. Third, body size (e.g., length or height) estimations can more easily be automated,

using e.g., image analysis software (Paavo *et al.*, 2008; Mallard, Le Bourlot & Tully, 2013), to rapidly process multiple individuals at a time.

In benthic ecology, ash-free dry mass (hereafter ‘AFDM’, in the older literature sometimes called ‘ash-free dry weight’ or simply ‘AFDW’) is often regarded as the most accurate predictor of macrofauna biomass, as it only includes biologically active tissue. Since AFDM estimations require the incineration of dried samples in a furnace at high temperature, adding considerable time and costs to analyses, many studies have reported how AFDM scales with estimations of wet- and dry mass, usually in the form of simple ratios as ‘conversion factors’ (e.g., AFDM/DM, in %) (Rumohr, Brey & Ankar, 1987; Ricciardi & Bourget, 1998). However, these ratios typically ignore the possibility that the relative mass of biologically active vs. non-active support tissue (e.g., protective exoskeleton or shell)—and therefore the AFDM/DM ratio—may change with macrofauna body size, as previously shown for disparate taxa like spiders (Andersen, 1979), vertebrates (Miller & Birchard, 2005) and trees (Niklas, 1995). This issue is important not only for obtaining accurate biomass conversions and estimations, but also for understanding how organismal investment in one type of structure may limit or constrain investment in other structures across ontogenetic development stages (Lease & Wolf, 2010).

Here we estimate and report relationships between body size, dry mass and ash-free dry mass for 14 of the most common aquatic, epibenthic invertebrate taxa found in shallow, vegetated habitats of the central Baltic Sea; one of the largest brackish water bodies on Earth. For each taxon we also assess whether the ash-free dry mass/dry mass ratio changes with body size. Our aim is to provide simple yet reliable size-based relationships that can be used to rapidly estimate organism body mass and, ultimately, biomass per sample.

## METHODS

### Study area

The Baltic Sea is a 415,000 km<sup>2</sup> large marginal sea situated in northern Europe (53–66°N; 10–30°E). A main feature is the presence of strong horizontal and vertical gradients in salinity, temperature and oxygen, that also undergo considerable temporal (e.g., seasonal) fluctuations (Voipio, 1981). The Baltic Sea is evolutionary very young (ca 6,000 years), and the shallow coastal areas have since the last glaciation been colonized by a mixture of marine, freshwater and brackish organisms, including crustaceans, gastropods, bivalves, polychaetes, hirudineans, nemerteans and insect larvae (Hansen, Wikström & Kautsky, 2008). As many marine and freshwater organisms in the Baltic Sea live near their physiological tolerance limits, they grow slower and smaller than in their original environment; e.g., the blue mussel *Mytilus edulis* (Tedengren & Kautsky, 1986). As a consequence, their size ranges—but also size:mass relationships and, potentially, AFDM/DM ratios—could differ from those reported for conspecifics in marine or freshwater areas (Rumohr, Brey & Ankar, 1987). An estimate of the effect of salinity on size:mass or DM:AFDM relationship was beyond the scope of our study, but our results could be compared to relationships in marine populations of the same taxa, if sampled and measured in the same way.

## Field sampling

During summer (May–Aug) 2014 we collected aquatic invertebrate macrofauna (>1 mm) in 32 shallow bays situated along a 360 km stretch of the central, Swedish Baltic Sea coastline (Fig. 1). The salinity in the area is generally low (ca. 5–7 psu) but fluctuates strongly with freshwater runoff and upwelling events. In each bay, a snorkeler sampled submerged aquatic vegetation and epibenthic macrofauna in 3–8 randomly selected stations (>30 m apart), by gently placing a 20 × 20 cm frame (with a 1 mm-mesh bag attached) on the sea bed, and collecting all organisms (aquatic vegetation and associated invertebrates) found above or on top of the sediment surface. The bag content was immediately transferred to a plastic bag, which was kept cold on ice until frozen (−20 °C), in most cases within 1–3 h.

## Body size estimations

Following thawing in room temperature, we identified intact invertebrate organisms to the highest taxonomic resolution feasible using standard literature. For the 14 most common taxa we then selected and measured the body size of 12–459 individuals per taxa (3,220 individuals in total), chosen to capture the full range of body sizes found across the 32 bays. The taxa included five gastropods (*Theodoxus fluviatilis*, *Hydrobia* spp., *Radix balthica*, *Potamopyrgus antipodarum*, *Bithynia tentaculata*), three bivalves (*Mytilus edulis*, *Limecola* (*Macoma*) *balthica* and *Cardidae* spp. (numerically dominated by *Parvicardium hauniense*), three crustaceans (*Amphibalanus improvisus*, *Idotea* spp., *Gammarus* spp.) and three insects (larval stages of Chironomidae spp., *Agraylea* spp. and *Phryganeidae* spp.) (see also Table 1). Body size (to the nearest 1 mm) was measured (based on standard procedures; Hayward & Ryland, 1995) as; (i) gastropod height along the central shell axis, (ii) bivalve length from anterior to posterior side, (iii) total length of *Gammarus* and *Idotea* spp. from tip of rostrum to last urosome, (iv) body width for *Amphibalanus improvisus*, and (v) total length of insect larvae from end of head to last segment. A higher size accuracy is definitely possible (e.g., to 0.1 or 0.01 mm using calipers or stereo lenses), but as most studies utilizing this type of data (including ours) will depend on 1000s of length measurements, the accuracy chosen was a realistic trade-off between time and precision.

## Estimations of dry- and ash-free dry mass

Following size estimations, the measured individuals were transferred to pre-dried and -weighed (nearest 0.0001 g) porcelain crucibles. For most size classes (except for very large and rare individuals), multiple individuals were typically pooled into the same crucible. This step underestimates actual variability in body mass between individuals, but was necessary as the low individual body masses (particularly AFDM) were near or below the reliable detection limit of the scale. We included multiple estimations of the same sizes, so that the number of biomass estimations (*N*) ranged from 10 to 42 per taxa. Samples were then dried at 60 °C for >48 h (until constant mass), and cooled to room temperature in a desiccator before weighing. To estimate ash-free dry mass, the crucibles were then transferred to a muffle furnace, incinerated (550 °C for 3 h), cooled and weighed again. Ash-free dry mass was calculated as dry mass minus ash mass.

**Table 1** Results of regression analyses estimating (i) the non-linear power relationship between body size and dry mass (DM) and (ii) ash-free dry mass (AFDM), (iii) the mean  $\pm$  1 SE AFDM/DM ratio (in %), and (iv) the linear relationship between body size and AFDM/DM ratio (in %), for 14 macroinvertebrate taxa in shallow coastal areas of the Baltic Sea. Letters within parentheses after taxa names denote classes.

Taxon	N	Body size vs. DM			Body size vs. AFDM			AFDM/DM	Body size vs. AFDM/DM		
		$\alpha \pm$ SE	$\beta \pm$ SE	$R^2$	$\alpha \pm$ SE	$\beta \pm$ SE	$R^2$	Mean % $\pm$ 1 SE	Intercept $\pm$ SE	Slope $\pm$ SE	$R^2$
<i>Bithynia tentaculata</i> L. (G)	25	0.598 $\pm$ 0.484 <sup>ns</sup>	2.117 $\pm$ 0.351 <sup>***</sup>	0.847	0.479 $\pm$ 0.511 <sup>ns</sup>	1.36 $\pm$ 0.472 <sup>**</sup>	0.668	19.133 $\pm$ 2.207	33.162 $\pm$ 3.878 <sup>***</sup>	−1.91 $\pm$ 0.452 <sup>+</sup>	0.424
<i>Hydrobia</i> spp. (G)	24	0.239 $\pm$ 0.041 <sup>***</sup>	2.134 $\pm$ 0.095 <sup>***</sup>	0.952	0.079 $\pm$ 0.029 <sup>+</sup>	1.441 $\pm$ 0.22 <sup>***</sup>	0.758	13.737 $\pm$ 1.155	19.791 $\pm$ 2.855 <sup>***</sup>	−0.633 $\pm$ 0.715 <sup>+</sup>	0.155
<i>Potamopyrgus antipodarum</i> Gray (G)	17	0.479 $\pm$ 0.511 <sup>ns</sup>	1.360 $\pm$ 0.472 <sup>**</sup>	0.919	0.021 $\pm$ 0.012 <sup>ns</sup>	2.447 $\pm$ 0.395 <sup>***</sup>	0.898	16.051 $\pm$ 1.399	6.063 $\pm$ 4.616 <sup>ns</sup>	2.653 $\pm$ 1.180 <sup>+</sup>	0.202
<i>Radix balthica</i> L. (G)	20	0.137 $\pm$ 0.035 <sup>***</sup>	2.355 $\pm$ 0.115 <sup>***</sup>	0.956	0.046 $\pm$ 0.018 <sup>+</sup>	2.119 $\pm$ 0.177 <sup>***</sup>	0.906	27.087 $\pm$ 2.233	35.338 $\pm$ 3.558 <sup>***</sup>	−1.794 $\pm$ 0.650 <sup>+</sup>	0.258
<i>Theodoxus fluviatilis</i> L. (G)	29	0.221 $\pm$ 0.065 <sup>***</sup>	2.683 $\pm$ 0.148 <sup>***</sup>	0.9492	0.015 $\pm$ 0.006 <sup>+</sup>	2.915 $\pm$ 0.194 <sup>***</sup>	0.912	13.044 $\pm$ 1.083	18.52 $\pm$ 2.396 <sup>***</sup>	−0.242 $\pm$ 0.494 <sup>+</sup>	0.159
<i>Cardidae</i> spp. (B)	33	0.134 $\pm$ 0.094 <sup>ns</sup>	2.848 $\pm$ 0.347 <sup>***</sup>	0.924	0.014 $\pm$ 0.013 <sup>ns</sup>	2.806 $\pm$ 0.486 <sup>***</sup>	0.879	12.358 $\pm$ 0.852	18.075 $\pm$ 1.468 <sup>***</sup>	−0.429 $\pm$ 0.325 <sup>+</sup>	0.364
<i>Limecola balthica</i> L. (B)	18	0.069 $\pm$ 0.024 <sup>+</sup>	2.820 $\pm$ 0.134 <sup>***</sup>	0.991	0.001 $\pm$ 0.002 <sup>ns</sup>	3.479 $\pm$ 0.673 <sup>***</sup>	0.92	12.717 $\pm$ 1.934	21.429 $\pm$ 2.98 <sup>***</sup>	−0.264 $\pm$ 0.372 <sup>+</sup>	0.383
<i>Mytilus edulis</i> L. (B)	24	0.030 $\pm$ 0.015 <sup>+</sup>	2.933 $\pm$ 0.153 <sup>***</sup>	0.991	0.006 $\pm$ 0.003 <sup>+</sup>	2.844 $\pm$ 0.147 <sup>***</sup>	0.978	14.189 $\pm$ 0.504	13.162 $\pm$ 1.044 <sup>***</sup>	0.078 $\pm$ 0.069 <sup>ns</sup>	0.011
<i>Amphibalanus improvisus</i> Darwin (C)	13	0.314 $\pm$ 0.205 <sup>ns</sup>	2.515 $\pm$ 0.289 <sup>***</sup>	0.976	0.036 $\pm$ 0.022 <sup>ns</sup>	2.289 $\pm$ 0.276 <sup>***</sup>	0.961	8.939 $\pm$ 0.550	11.044 $\pm$ 1.064 <sup>***</sup>	−0.397 $\pm$ 0.179 <sup>+</sup>	0.246
<i>Gammarus</i> spp. (C)	37	0.047 $\pm$ 0.032 <sup>ns</sup>	2.111 $\pm$ 0.265 <sup>***</sup>	0.926	0.033 $\pm$ 0.028 <sup>ns</sup>	2.05 $\pm$ 0.32 <sup>***</sup>	0.863	58.966 $\pm$ 1.519	63.062 $\pm$ 2.616 <sup>***</sup>	−0.389 $\pm$ 0.307 <sup>ns</sup>	0.017
<i>Idothea</i> spp. (C)	42	0.001 $\pm$ 0.001 <sup>ns</sup>	3.592 $\pm$ 0.200 <sup>***</sup>	0.949	0.001 $\pm$ 0.001 <sup>ns</sup>	3.850 $\pm$ 0.249 <sup>***</sup>	0.919	61.505 $\pm$ 1.659	66.183 $\pm$ 3.457 <sup>***</sup>	−0.550 $\pm$ 0.358 <sup>ns</sup>	0.032
<i>Agraylea</i> spp. (I)	13	0.001 $\pm$ 0.002 <sup>ns</sup>	3.410 $\pm$ 0.721 <sup>**</sup>	0.820	0.001 $\pm$ 0.002 <sup>ns</sup>	3.432 $\pm$ 0.769 <sup>***</sup>	0.833	85.967 $\pm$ 3.769	88.893 $\pm$ 7.725 <sup>***</sup>	0.570 $\pm$ 1.277 <sup>ns</sup>	−0.097
<i>Chironomidae</i> spp. (I)	38	0.014 $\pm$ 0.016 <sup>ns</sup>	1.383 $\pm$ 0.290 <sup>***</sup>	0.600	0.008 $\pm$ 0.006 <sup>ns</sup>	1.544 $\pm$ 0.321 <sup>***</sup>	0.533	79.307 $\pm$ 2.643	78.633 $\pm$ 6.947 <sup>***</sup>	0.070 $\pm$ 0.688 <sup>ns</sup>	−0.027
<i>Phryganeidae</i> spp. (I)	10	0.001 $\pm$ 0.001 <sup>ns</sup>	3.176 $\pm$ 0.649 <sup>***</sup>	0.746	0.001 $\pm$ 0.001 <sup>ns</sup>	3.207 $\pm$ 0.611 <sup>***</sup>	0.789	91.851 $\pm$ 2.137	86.64 $\pm$ 3.558 <sup>***</sup>	0.382 $\pm$ 0.185 <sup>ns</sup>	0.290

**Notes.**

G, Gastropoda; B, Bivalvia; C, Crustacea; I, Insecta (larvae);  $\alpha$  and  $\beta$ , normalization and scaling constant for power equations, respectively.


<sup>ns</sup>  $p < 0.05$ .

<sup>+</sup>  $p < 0.05$ .

<sup>\*\*</sup>  $p < 0.01$ .

<sup>\*\*\*</sup>  $p < 0.001$ .

Values in bold mark those significant (at  $\alpha = 0.05$ ). Note:  $R^2$  were derived from linear log–log models.



**Figure 1** Maps of Scandinavia (small image) and the sampling area. Filled circles mark the position of the 32 sampled bays. Numbers along the  $x$ - and  $y$ -axis are longitude and latitude, respectively.



## Statistical analyses

We estimated taxon-specific body size:body mass relationship using non-linear regression in the form of the power equation:

$$\text{body mass} = \alpha \times \text{size}^{\beta}$$

where *body mass* is the individual mass (mg DM or AFDM), *size* is the body size (length/height, in mm),  $\alpha$  is a normalization constant, and  $\beta$  is the scaling constant. Body mass typically scales with size in a power relationship, and initial data exploration showed that power equations provided a superior fit compared to linear, log or exponential relationships. As regression coefficients ( $R^2$ ) are an inadequate measure of fit for non-linear regression models (Spiess & Neumeyer, 2010), we report SE for  $\alpha$  and  $\beta$ . However, for the sake of simplicity we also estimated the linear log-log relationship between body size and biomass, and report the  $R^2$  for those models (see e.g., Lease & Wolf, 2010).

For each taxon we also calculated the mean ( $\pm 1$  SE) AFDM/DM ratio (in %); a commonly used conversion factor in macroinvertebrate studies (see e.g., Ricciardi & Bourget, 1998). We then used linear regression to test whether body size (in mm) affected the AFDM/DM ratio. Prior to analyses we checked assumptions of normality (by plotting predicted vs. observed quantiles) and homoscedasticity (by plotting predicted vs. observed residuals). All analyses were conducted in R v. 3.2.3 (R Core Team, 2016).

## RESULTS

### Relationships between body size and individual biomass


The relationships between body size (mm), individual dry mass (mg DM) and ash-free dry mass (mg AFDM) for all 14 taxa are displayed in Figs. 2A–2H, and the parameters (and their fit) are presented in Table 1. For most of the taxa, body size was a very good predictor of individual DM, as demonstrated by low SE and  $R^2$  near 1. The model fits were slightly poorer for the three insect taxa ( $R^2 = 0.60$ – $0.82$ ) and the gastropod *Bithynia tentaculata* ( $R^2 = 0.85$ ) than for the other ten taxa. For a majority (12 out of 14) of the taxa, the scaling constants ( $\beta$ ) were well above 2 (2.110–3.590). The exceptions were the small gastropod *Potamopyrgus antipodarum* and chironomid larvae, which had constants closer to 1 ( $\beta = 1.368$  and 1.383, respectively).

Body size was also a very good predictor of AFDM, even though model fits (based on SE and  $R^2$ ) were slightly poorer than for DM (Table 1). Just as for DM relationships, the model fits (based on SE and  $R^2$ ) were best for gastropods, molluscs and crustaceans. The scaling constants ( $\beta$ ) were for most taxa quite similar to those reported for the DM relationships, with the exception of a higher constant for *P. antipodarum* ( $\beta = 2.447$ ) and a lower constant for *Bithynia tentaculata* ( $\beta = 1.360$ ).

### Influence of organism body size on AFDM/DM ratios

The AFDM/DM ratios (mean %  $\pm$  SE) per taxa are also presented in Table 1. As expected, there were consistent differences between the four major taxonomic groups studied, with low AFDM content in bivalves and gastropods (12–27%), who's calcium carbonate shell





**Figure 2** Best-fitting relationships between body size (length or height, see ‘Methods’) and (A–D) dry mass (mg. DM), (E–H) ash-free dry mass (mg. AFDM) and (I–L) AFDM/DM ratio (% AFDM), for 14 taxa—five gastropods, three bivalves, three crustaceans and three insect larvae—sampled in coastal areas of the central Baltic Sea. For model parameters and estimates of fit, see [Table 1](#).

makes up the major part of whole-body biomass, to higher AFDM content in chitin-shelled crustaceans (ca 60%), and the highest content in insect larvae (86–92%).

Results of simple linear regression showed that for more than half (8 out of 14) of the taxa surveyed, body size clearly affected the AFDM/DM ratio ([Table 1](#) and [Figs. 2I–2L](#)). For four out of five gastropods, two out of three bivalves, as well as the sessile, calcite-shelled crustacean *Amphibalanus improvisus*, the AFDM/DM ratio decreased linearly with body size. For the small gastropod *Potamopyrgus antipodarum* body size instead had a positive influence

on AFDM/DM. However, the *P. antipodarum* size range was very narrow (2–4 mm) and the intercept was not different from 0 (Table 1), suggesting a relatively poor model fit. Moreover, there was no size effect found in the blue mussel *Mytilus edulis* (Table 1). Finally, in contrast to the size effects found for most of the hard-shelled molluscs, there was no influence of body size on AFDM/DM in any of the chitin-shelled crustaceans or insect larvae (Table 1 and Figs. 2I–2L).

## DISCUSSION

Estimating organism biomass is one of the most common, important but also resource-consuming tasks in ecological work, particularly when it comes to small-bodied, highly abundant and diverse macroscopic invertebrates. Many previous studies have shown that more easily measured variables like invertebrate wet (fresh) mass (e.g., Ricciardi & Bourget, 1998) or body size (e.g., Smock, 1980) can be used as proxies to reliably predict both the dry- and ash-free dry body mass, thereby simplifying and speeding up biomass estimations. Here, we first complement this literature by reporting how body mass scales with body size for 14 of the most common epibenthic invertebrate taxa found in shallow coastal areas of the Baltic Sea. Moreover, we demonstrate that for a majority of the studied molluscs, the ratio between organism dry- and ash-free dry mass—an often-used conversion factor (e.g., Rumohr, Brey & Ankar, 1987; Ricciardi & Bourget, 1998)—decreases predictably with body size. Thus, our results can be used to quickly estimate the biologically active biomass of individual organisms based on their size, and when combined with density data, accurately estimate biomass per unit area.

### Body size as a proxy for dry- and ash-free dry mass

For a majority of the studied taxa, body size was a good predictor of both dry mass and ash-free dry mass. The model fits were slightly poorer for ash-free dry mass (AFDM); most likely a consequence of the fact that even though multiple individuals of the same size were pooled, the low individual AFDM of many organisms (in the vicinity of 1 mg) challenged the accuracy of the scale. Comparisons between the 14 taxa studied (Table 1) show that particularly within the gastropods and crustaceans, the scaling ( $\beta$ ) constants differ quite substantially between taxa (see the different slopes in Fig. 2 and  $\beta$  coefficients in Table 1). These differences emphasize the need for taxon-specific relationships to accurately predict biomass, and the potential dangers in either ignoring body size or substituting relationships between taxa. Consequently, our power equations (Table 1) can be used in a simple yet reliable way to estimate organism dry- or ash-free dry mass based on standard body size measurements. Future studies should ideally also assess how these relationships vary in time and space (e.g., over seasons), for even more accurate biomass estimations. Size-based biomass estimations are likely to speed up laboratory work considerably; for example, Casagrande & Boudouresque (2002) showed that sieve-based size estimations speeded up estimations of body biomass of the gastropod *Hydrobia ventricosa* by 20–30 times. Consequently, our size-based estimations of invertebrate biomass are likely to free up considerable work resources (time, man-power, money) that can be used to e.g., collect and process more samples.

## The influence of body size on AFDM/DM ratios

For most of the taxa with a calcium-carbonate (molluscs) or calcite shell (the barnacle *Amphibalanus improvisus*), we found a significant negative influence of body size on the AFDM/DM ratio; a commonly reported and often-used conversion factor in macrofauna studies (e.g., [Rumohr, Brey & Ankar, 1987](#); [Ricciardi & Bourget, 1998](#)). In other words, the proportional mass of biologically active vs. non-active tissue (shell, hard mouth parts, etc.) decreased with body size. There are at least two possible and complementary explanations for this relationship. First, while the rate of growth in length of mollusc shells typically decreases with age, new shell layers are consistently added on a yearly basis ([Negus, 1966](#)). This results in increasingly thicker, and therefore disproportionately heavier, shells with mussel length, and a higher shell:tissue mass ratio. Second, our sampling was conducted during summer; a season when a majority of adult molluscs (here represented by the larger individuals per taxa) most likely had spawned and temporarily lost a considerable proportion of their biologically active tissue ([Kautsky, 1982](#)). The slopes of the significant regressions ([Table 1](#), median =  $-1.26$ ) suggest that failing to incorporate the potential influence of body size can strongly reduce the accuracy of AFDM estimations based on dry mass (and presumably also wet mass), particularly if there is considerable variability in body size in the samples. The somewhat surprising lack of size influence in the common blue mussel *Mytilus edulis* was not investigated in detail, but could be caused by (i) the lack of small shell-crushing mussel predators in the area (e.g., crabs), who otherwise are known to trigger thicker mussel shells ([Freeman, 2007](#)), and/or (ii) the relatively low salinity, which causes the small, osmotically stressed *M. edulis* to invest considerably more energy into osmosis and soft tissue production, than in thicker shells ([Kautsky, Johannesson & Tedengren, 1990](#)).

In contrast to the results for molluscs, there was no size effect on AFDM/DM ratios for the chitin-shelled insects and crustaceans. These results fit well with those reported in previous studies, for example of terrestrial insects, for which exoskeletal chitin scales isometrically (1:1) with body size ([Lease & Wolf, 2010](#)). In summary, our results suggest that body size can play an important but hitherto underestimated role when estimating organism AFDM based on dry (and possibly, wet) mass, particularly for shelled molluscs.

## CONCLUSIONS

Using samples of epibenthic macroinvertebrates collected in 32 shallow bays along a 360 km stretch of the Swedish Baltic Sea coast, we show that for 14 of the most common macrofauna taxa, organism body size scales predictably with individual dry mass and ash-free dry mass in the form of power relations. The good model fits suggest the taxon-specific equations reported here can be used to predict individual biomass based on organism size, thereby speeding up estimations of macrofauna biomass. Moreover, for the vast majority of the molluscs studied, we find a negative relationship between body size and AFDM/DM ratio; a commonly used conversion factor in macrofauna studies. Consequently, future studies utilizing AFDM/DM ratios should carefully assess the potential influence of body size and spatial-temporal variability, to ensure reliable biomass estimations.

## ACKNOWLEDGEMENTS

We acknowledge the field and/or laboratory assistance of (in alphabetic order) T Amgren, E Anderberg, F Ek, P Jacobsson, G Johansson, C Jönander, L Näslund, O Pettersson, M van Regteren, S Skoglund, E Svartgren, V Thunell, C Åkerlund and M Åkerman. We thank the water right owners around each study bay for facilitating the field sampling. This study is a product of project Plant-Fish ([www.plantfish.se](http://www.plantfish.se)).

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This study was funded by Formas (grant 2013-1074), His Majesty Carl XVI Gustaf's Foundation for Science and Education (grant HMK-2014-0002), the Baltic Sea 2020 foundation (project "Levande kust"), Stockholm University Baltic Sea Centre (Askö grants and support to MSc students), and in-kind support from Groningen University (Netherlands) to BDHK Eriksson. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Formas: 2013-1074.

His Majesty Carl XVI Gustaf's Foundation for Science and Education: HMK-2014-0002.  
Baltic Sea 2020 foundation.

Stockholm University Baltic Sea Centre.

Groningen University.

### Competing Interests

Göran Sundblad is an employee of AquaBiota Water Research. His work in relation to this study was fully funded by a grant from Formas, the Swedish research council for sustainable development.

### Author Contributions

- Johan Eklöf conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables.
- Åsa Austin, Serena Donadi, Britas D.H.K. Eriksson and Göran Sundblad performed the experiments, contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- Ulf Bergström and Joakim Hansen performed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, reviewed drafts of the paper.

### Data Availability

The following information was supplied regarding data availability:

The raw data has been supplied as [Data S1](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.2906#supplemental-information>.

## REFERENCES

- Andersen SO. 1979. Biochemistry of insect cuticle. *Annual Review of Entomology* 24:29–59 DOI 10.1146/annurev.en.24.010179.000333.
- Brey T, Muller-Wiegmann C, Zittier ZMC, Hagen W. 2010. Body composition in aquatic organisms—a global data bank of relationships between mass, elemental composition and energy content. *Journal of Sea Research* 64(3):334–340 DOI 10.1016/j.seares.2010.05.002.
- Brey T, Rumohr H, Ankar S. 1988. Energy content of macrobenthic invertebrates: general conversion factors from weight to energy. *Journal of Experimental Marine Biology and Ecology* 117(3):271–278 DOI 10.1016/0022-0981(88)90062-7.
- Casagrande C, Boudouresque CF. 2002. A sieving method for rapid determination of size-frequency distribution of small gastropods. Example of the mud snail *Hydrobia ventrosa*. *Hydrobiologia* 485(1):143–152 DOI 10.1023/A:1021371308753.
- Edgar GJ. 1990. The use of the size structure of benthic macrofaunal communities to estimate faunal biomass and secondary production. *Journal of Experimental Marine Biology and Ecology* 137(3):195–214 DOI 10.1016/0022-0981(90)90185-F.
- Enquist BJ, Niklas KJ. 2001. Invariant scaling relations across tree-dominated communities. *Nature* 410:655–660 DOI 10.1038/35070500.
- Freeman AS. 2007. Specificity of induced defenses in *Mytilus edulis* and asymmetrical predator deterrence. *Marine Ecology Progress Series* 334:145–153 DOI 10.3354/meps334145.
- Frithsen JB, Rudnick DT, Doering PH. 1986. The determination of fresh organic carbon weight from formaldehyde preserved macrofaunal samples. *Hydrobiologia* 133(3):203–208 DOI 10.1007/BF00005591.
- Gruner DS, Smith JE, Seabloom EW, Sandin SA, Ngai JT, Hillebrand H, Harpole WS, Elser JJ, Cleland EE, Bracken MES, Borer ET, Bolker BM. 2008. A cross-system synthesis of consumer and nutrient resource control on producer biomass. *Ecology Letters* 11(7):740–755 DOI 10.1111/j.1461-0248.2008.01192.x.
- Hansen JP, Wikström SA, Kautsky L. 2008. Effects of water exchange and vegetation on the macroinvertebrate fauna composition of shallow land-uplift bays in the Baltic Sea. *Estuarine, Coastal and Shelf Science* 77(3):535–547 DOI 10.1016/j.ecss.2007.10.013.
- Hayward PJ, Ryland JS. 1995. *Handbook of the marine fauna of North-West Europe*. New York: Oxford University Press, 800 pp.
- Hjörleifsson E, Klein-MacPhee G. 1992. Estimation of live standard length of winter flounder *Pleuronectes americanus* larvae from formalin-preserved, ethanol-preserved and frozen specimens. *Marine Ecology Progress Series* 82:13–19 DOI 10.3354/meps082013.

- Howmiller RP. 1972.** Effects of preservatives on weights of some common macrobenthic invertebrates. *Transactions of the American Fisheries Society* **101**(4):743–746 DOI [10.1577/1548-8659\(1972\)101<743:EOPOWO>2.0.CO;2](https://doi.org/10.1577/1548-8659(1972)101<743:EOPOWO>2.0.CO;2).
- Kapiris K, Miliou H, Moraitou-Apostolopoulou M. 1997.** Effects of formaldehyde preservation on biometrical characters, biomass and biochemical composition of *Acartia clausi* (Copepoda, Calanoida). *Helgolander Meeresuntersuchungen* **51**(1):95–106 DOI [10.1007/BF02908757](https://doi.org/10.1007/BF02908757).
- Kautsky N. 1982.** Quantitative studies on gonad cycle, fecundity, reproductive output and recruitment in a baltic *Mytilus edulis* population. *Marine Biology* **68**(2):143–160 DOI [10.1007/BF00397601](https://doi.org/10.1007/BF00397601).
- Kautsky N, Johannesson K, Tedengren M. 1990.** Genotypic and phenotypic differences between Baltic and North Sea populations of *Mytilus edulis* evaluated through reciprocal transplantations. I. Growth and morphology. *Marine Ecology Progress Series* **59**:203–210 DOI [10.3354/meps059203](https://doi.org/10.3354/meps059203).
- Lease HM, Wolf BO. 2010.** Exoskeletal chitin scales isometrically with body size in terrestrial insects. *Journal of Morphology* **271**(6):759–768 DOI [10.1002/jmor.10835](https://doi.org/10.1002/jmor.10835).
- Leuven RSEW, Brock TCM, Van Druten HAM. 1985.** Effects of preservation on dry- and ash-free dry weight biomass of some common aquatic macro-invertebrates. *Hydrobiologia* **127**(2):151–159 DOI [10.1007/BF00004193](https://doi.org/10.1007/BF00004193).
- Mallard F, Le Bourlot V, Tully T. 2013.** An automated image analysis system to measure and count organisms in laboratory microcosms. *PLOS ONE* **8**(5):e64387 DOI [10.1371/journal.pone.0064387](https://doi.org/10.1371/journal.pone.0064387).
- Mason WT, Lewis PA, Weber CI. 1983.** An evaluation of benthic macroinvertebrate biomass methodology. *Environmental Monitoring and Assessment* **3**(1):29–44 DOI [10.1007/BF00394030](https://doi.org/10.1007/BF00394030).
- Miller K, Birchard GF. 2005.** Influence of body size on shell mass in the ornate box turtle, terrapene ornata. *Journal of Herpetology* **39**(1):158–161 DOI [10.1670/0022-1511\(2005\)039\[0158:IOBSOS\]2.0.CO;2](https://doi.org/10.1670/0022-1511(2005)039[0158:IOBSOS]2.0.CO;2).
- Negus C. 1966.** A quantitative study of growth and production of unionid mussels in the River Thames at reading. *Journal of Animal Ecology* **35**:513–532 DOI [10.2307/2489](https://doi.org/10.2307/2489).
- Niklas KJ. 1995.** Size-dependent allometry of tree height, diameter and Truck-taper. *Annals of Botany* **75**:217–227 DOI [10.1006/anbo.1995.1015](https://doi.org/10.1006/anbo.1995.1015).
- Paavo B, Ziegelmeyer A, Lavric E, Probert PK. 2008.** Morphometric correlations and body mass regressions for *Armandia maculata*, *Aglaophamus macroura* (Polychaeta), and *Zethalia zelandica* (Gastropoda). *New Zealand Journal of Marine and Freshwater Research* **42**(1):85–91 DOI [10.1080/00288330809509938](https://doi.org/10.1080/00288330809509938).
- Perez-Harguindeguy N, Diaz S, Garnier E, Lavorel S, Poorter H, Jaureguiberry P, Bret-Harte MS, Cornwell WK, Craine JM, Gurvich DE, Urcelay C, Veneklaas EJ, Reich PB, Poorter L, Wright IJ, Ray P, Enrico L, Pausas JG, De Vos AC, Buchmann N, Funes G, Quetier F, Hodgson JG, Thompson K, Morgan HD, Ter Steege H, Van der Heijden MGA, Sack L, Blonder B, Poschlod P, Vaieretti MV, Conti G, Staver AC, Aquino S, Cornelissen JHC. 2013.** New handbook for standardised measurement

- of plant functional traits worldwide. *Australian Journal of Botany* **61**:167–234 DOI 10.1071/BT12225.
- R Core Team.** 2016. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <https://www.r-project.org/>.
- Ricciardi A, Bourget E.** 1998. Weight-to-weight conversion factors for marine benthic macroinvertebrates. *Marine Ecology-Progress Series* **163**:245–251 DOI 10.3354/meps163245.
- Rumohr H, Brey T, Ankar S.** 1987. A compilation of biometric conversion factors for benthic invertebrates of the Baltic Sea. *The Baltic Marine Biologists* **9**:1–56.
- Sabo JL, Bastow JL, Power ME.** 2002. Length–mass relationships for adult aquatic and terrestrial invertebrates in a California watershed. *Journal of the North American Benthological Society* **21**(2):336–343 DOI 10.2307/1468420.
- Smock LA.** 1980. Relationships between body size and biomass of aquatic insects. *Freshwater Biology* **10**(4):375–383 DOI 10.1111/j.1365-2427.1980.tb01211.x.
- Spiess A-N, Neumeyer N.** 2010. An evaluation of R<sup>2</sup> as an inadequate measure for nonlinear models in pharmacological and biochemical research: a Monte Carlo approach. *BMC Pharmacology* **10**:6 DOI 10.1186/1471-2210-10-6.
- Tedengren M, Kautsky N.** 1986. Comparative study of the physiology and its probable effect on size in Blue Mussels (*Mytilus edulis* L.) from the North Sea and the Northern Baltic Proper. *Ophelia* **25**(3):147–155 DOI 10.1080/00785326.1986.10429746.
- Voipio A.** 1981. *The Baltic sea*. 1st edition. Vol. 30. Amsterdam: Elsevier Science.
- Widbom B.** 1984. Determination of average individual dry weights and ash-free dry weights in different sieve fractions of marine meiofauna. *Marine Biology* **84**(1):101–108 DOI 10.1007/BF00394532.